

ACTIVITY RESERVOIRS, REPEATING AND NON-REPEATING ACTIVITY OF COMET 67P.

N. Oklay¹ and The OSIRIS Team ¹DLR Institute for Planetary Research, Rutherford Str. 2, 12489 Berlin, Germany (Nilda.OklayVincent @dlr.de)

Long-term high spatial resolution observations of comet 67P during the Rosetta mission allowed us to study cometary activity in details. In some cases, direct connection of the events to the nucleus could be observed ([1], [2], [3]), in others the activity sources could be identified via indirect methods ([4], [5], [6], [7], [8], [9]). Systematic cataloguing of activity events observed during the mission revealed the existence of several activity source concentrations ([8], [9]). Not surprisingly, all those sources were concentrated on long-lasting H₂O ice rich areas ([1], [4], [8], [10]). By investigating surface modifications and spectral variations of those areas, we are able to estimate the local subsurface structures ([11]). Once we combine all available information from various Rosetta instruments for specific events, we can also estimate local volatile and supervolatile contents.

Majority of the activity events observed during Rosetta mission were repeating at different time-scales. Jet like collimated dust features observed on the limb were following the direct solar energy input, therefore, switched on/off diurnally. However, numerous jet like features observed on the surface were repeating but not diurnally ([11]). Additional to those, there were non-periodically repeating outbursts from the same sources but displaying various plume morphologies.

In this presentation, I will first summarize properties of the activity events observed during the Rosetta mission, and then focus on studies of several activity sources.

References:

- [1] Oklay et al., 2016, A&A: ‘Variation of comet 67P/Churyumov-Gerasimenko in regions showing activity’
- [2] Vincent et al., 2015, Nature, ‘Large heterogeneities in comet 67P as revealed by active pits from sinkhole collapse’
- [3] Agarwal et al., 2017, MNRAS, ‘Evidence of sub-surface energy storage in comet 67P from the outburst of 2016 July 03’
- [4] Vincent et al., 2016, A&A, ‘Are fractured cliffs the source of cometary dust jets? Insights from OSIRIS/Rosetta at 67P/Churyumov-Gerasimenko’
- [5] Vincent et al., 2016, MNRAS. ‘Summer fireworks on comet 67P’
- [6] Knollenberg et al., 2016, A&A: ‘A mini outburst from the nightside of comet 67P/Churyumov-Gerasimenko observed by the OSIRIS camera on Rosetta’
- [7] Shi et al., 2016, A&A: ‘Sunset jets observed on comet 67P/Churyumov-Gerasimenko sustained by subsurface thermal lag’
- [8] Lai et al., submitted, ‘Seasonal variations of the source regions of the dust jets on comet 67P/Churyumov-Gerasimenko’
- [9] Lin et al., in preparation
- [10] Oklay et al., 2017, MNRAS: ‘Long-term survival of surface water ice on comet 67P’
- [11] Oklay, 2018, LPSC: ‘Large sub-surface volatile reservoirs of comet 67P’