

Asymptotics for spherical particle motion in a spherically expanding flow.

V. V. Zakharov^{1,2}, S. L. Ivanovski^{2,3}, J.-F. Crifo⁴, V. Della Corte^{3,2}, A. Rotundi^{3,2}, M. Fulle⁵

- 1) Laboratoire de Meteorologie Dynamique, Universite Pierre et Marie Curie, 4 place Jussieu, 75252 Paris, France
- 2) INAF - Institute for Space Astrophysics and Planetology, Area Ricerca Tor Vergata, Via Fosso del Cavaliere I-100, Rome 00133, Italy
- 3) Università degli Studi di Napoli “Parthenope”, Dip. di Scienze e Tecnologie, CDN I-IC4, Naples 80143, Italy
- 4) LATMOS, CNRS/UVSQ/IPSL, 11 boulevard d'Alembert, 78280 Guyancourt, France
- 5) INAF - Osservatorio Astronomico di Trieste, Via Tiepolo I-11, Trieste 34143, Italy

In the context of an increasing number of complex multiparametric dust coma models, it was found convenient to construct an elementary model with a minimum number of parameters selected to represent the key processes acting on the dust. The model outputs can be used as a reference evaluation of these processes with rough estimates of the resulting dust properties, e.g. velocity. The present work introduces three universal and dimensionless parameters, which characterize the dust motion in an expanding flow and computes, as a function of these parameters: 1) the dust terminal velocity; 2) the time to acquire it; and 3) the distance at which it is acquired. The motion of dust grains is presented as a system of dimensionless ordinary differential equations, the solution of which depends upon the above mentioned three parameters. The numerical integration of this system was performed over a wide range of parameter space covering the whole range of physically possible conditions. Precomputed results of dust terminal velocity, time and distance at which it is reached are presented in dimensionless form. To obtain dimensional values for a particular case it is sufficient to perform algebraic operations.

In agreement with our model: 1) GIADA dust particle speed measurements are consistent with the calculated terminal velocities; 2) OSIRIS data constrain the dust acceleration limited within six nuclear radii for a broad range of particle sizes.