

Colours, spectral slopes and phase curves of 67P/C.G.'s nucleus as observed by Rosetta/OSIRIS

Feller, C.¹; Fornasier, S.¹; Hasselmann, P.¹; Barucci, MA.¹; Deshapriya, JDP.¹; Sierks, H.² and the OSIRIS Team

1: LESIA, Observatoire de Paris, Univ. PSL, CNRS, Univ. Paris Diderot, Sorbonne Paris Cité, Sorbonne Univ., 5 place Jules Janssen, 92195 Meudon, France; 2: Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg, 3, 37077, Goettingen, Germany;

During the 26 months the Rosetta probe followed the nucleus of comet 67P/Churyumov-Gerasimenko, the OSIRIS instrument surveyed extensively its surface at different spatial resolutions and under multiple phase angles. This allowed to characterize the overall surface of the nucleus. Furthermore the three flybys manoeuvres, that the spacecraft accomplished respectively in August 2014, February 2015 and April 2016 [7, 5, 9], most notably allowed to characterize the nucleus' surface at high resolution.

While the first flyby manoeuver was performed at an altitude of ~ 300 km, thus allowing the OSIRIS instrument to map most of the nucleus' surface, the other flybys, performed at ~ 10 and ~ 30 km over the big lobe of the comet and around the frontiers of the Imhotep depression, have permitted the OSIRIS instrument to study with a sub-meter resolution the spectrophotometric and photometric properties of the transition areas between different morphological units [3, 4].

The extensive amount of OSIRIS observations collected during those 26 months have also allowed to investigate the presence and resilience of areas locally enriched in water-ice material [11, 1, 2, 6, 10], as well as the evolution of the superficial layers' composition in between seasons and during sunrise [8].

We will summarize the findings of the OSIRIS composition team regarding the global spectrophotometric and photometric properties of the nucleus, as well as present the results of the photometric analysis of the April 2016 flyby observations and compare them with those obtained during the previous flybys.

References

- [1] M. A. Barucci, G. Filacchione, S. Fornasier, et al. Detection of exposed H₂O ice on the nucleus of comet 67P/Churyumov-Gerasimenko. as observed by Rosetta OSIRIS and VIRTIS instruments. *Astronomy and Astrophysics*, 595:A102, Nov. 2016.
- [2] J. D. P. Deshapriya, M. A. Barucci, S. Fornasier, et al. Spectrophotometry of the Khonsu region on the comet 67P/Churyumov-Gerasimenko using OSIRIS instrument images. *Monthly Notices of the Royal Astronomical Society*, 462:S274–S286, Nov. 2016.
- [3] M. R. El-Maarry, N. Thomas, L. Giacomini, et al. Regional surface morphology of comet 67P/Churyumov-Gerasimenko from Rosetta/OSIRIS images. *Astronomy and Astrophysics*, 583:A26, Nov. 2015.
- [4] M. R. El-Maarry, N. Thomas, A. Gracia-Berná, et al. Regional surface morphology of comet 67P/Churyumov-Gerasimenko from Rosetta/OSIRIS images: The southern hemisphere. *Astronomy and Astrophysics*, 593:A110, Sept. 2016.
- [5] C. Feller, S. Fornasier, P. H. Hasselmann, et al. Decimetre-scaled spectrophotometric properties of the nucleus of comet 67P/Churyumov-Gerasimenko from OSIRIS observations. *Monthly Notices of the Royal Astronomical Society*, 462:S287–S303, Nov. 2016.
- [6] S. Fornasier, C. Feller, J. C. Lee, et al. The highly active Anhur-Bes regions in the 67P/Churyumov - Gerasimenko comet: results from OSIRIS/ROSETTA observations. *MNRAS*, July 2017.
- [7] S. Fornasier, P. H. Hasselmann, M. A. Barucci, et al. Spectrophotometric properties of the nucleus of comet 67P/Churyumov-Gerasimenko from the OSIRIS instrument onboard the ROSETTA spacecraft. *Astronomy and Astrophysics*, 583:A30, Nov. 2015.
- [8] S. Fornasier, S. Mottola, H. U. Keller, et al. Rosetta's comet 67P/Churyumov-Gerasimenko sheds its dusty mantle to reveal its icy nature. *Science*, 354:1566–1570, Dec. 2016.
- [9] P. H. Hasselmann, M. A. Barucci, S. Fornasier, et al. The opposition effect of 67P/Churyumov-Gerasimenko on post-perihelion Rosetta images. *Monthly Notices of the Royal Astronomical Society*, 469:S550–S567, July 2017.
- [10] N. Oklay, S. Mottola, J.-B. Vincent, et al. Long-term survival of surface water ice on comet 67P. *Monthly Notices of the Royal Astronomical Society*, 469:S582–S597, July 2017.
- [11] A. Pommerol, B. Jost, O. Poch, et al. The SCITEAS experiment: Optical characterizations of sublimating icy planetary analogues. *Planetary Space Sciences*, 109:106–122, May 2015.