

Oxygen Isotopes in Water in the Coma of Comet 67P / Churyumov-Gerasimenko as measured with the Rosetta / ROSINA Double Focusing Mass Spectrometer

I. Schroeder (1), K. Altwegg (1,2), H. Balsiger (1), J. J. Berthelier (3), J. De Keyser (4), B. Fiethe (5), S. A. Fuselier (6), S. Gasc (1), T. I. Gombosi (7), M. Rubin (1), T. Sémon (1), C.-Y. Tzou (1), S. F. Wampfler (2), P. Wurz (1,2)

- (1) Physikalisches Institut, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland
- (2) Center for Space and Habitability, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland
- (3) LATMOS, 4 Avenue de Neptune, F-94100 Saint-Maur, France
- (4) Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Ringlaan 3, B-1180, Brussels, Belgium
- (5) Institute of Computer and Network Engineering (IDA), TU Braunschweig, Hans-Sommer-Strasse 66, D-38106 Braunschweig, Germany
- (6) Space Science Division, Southwest Research Institute, 6220 Culebra Road, San Antonio, TX 78228, USA
- (7) Department of Atmospheric, Oceanic and Space Sciences, University of Michigan, 2455 Hayward, Ann Arbor, MI 48109, USA

Abstract

Comets are widely considered to contain some of the most pristine material in the Solar System [1]. The degree of isotopic fractionation – the enrichment or depletion of an isotope in a molecule, relative to its initial abundance – observed in a comet is sensitive to the environmental conditions at the time of the comet's formation [2]. Therefore, measurements of isotopic abundances in cometary ices reveal important information regarding the early Solar System's composition, density, temperature and the amount of radiation present before the accretion of solid bodies, when the molecules were being formed during the chemical evolution of the presolar cloud to the protosolar nebula and protoplanetary disc. They are therefore vital to understanding and reconstructing the history and origins of material in the Solar System [3].

The $^{16}\text{O} / ^{18}\text{O}$ ratio of CO_2 in the coma of the comet 67P / Churyumov-Gerasimenko was previously measured by Hässig et al. (2016) [2] with the ESA spacecraft Rosetta's ROSINA instrument package's Double Focusing Mass Spectrometer (DFMS) and found to be 494 ± 8 , which is consistent within 1σ uncertainty with the value of 499 calculated by Lodders (2003) [4] for the solar system abundances, but not with the ratio of 530 ± 2 measured for the solar wind by McKeegan et al. (2011) [5].

In this study, the $^{16}\text{O} / ^{18}\text{O}$ ratio of H_2O in the coma of the comet 67P, as measured by the Rosetta / ROSINA DFMS, was found from the ratio of $\text{H}_2\text{O} / \text{H}_2^{18}\text{O}$ to be 445 ± 35 , which represents a 12% enrichment of ^{18}O compared with the terrestrial value [4] of 499 and would be consistent with the comet containing primordial water, in accordance with leading self-shielding models, which hypothesise primordial water to be between 5% to 20% more enriched in the heavier oxygen isotopes than terrestrial water [6].

Acknowledgements

ROSINA would not have produced such outstanding results without the work of the many engineers, technicians, and scientists involved in the mission, in the Rosetta spacecraft, and in the ROSINA instrument team over the last 20 years, whose contributions are gratefully acknowledged. Rosetta is a European Space Agency (ESA) mission with contributions from its member states and NASA. We acknowledge herewith the work of the whole ESA Rosetta team.

Work at University of Bern was funded by the State of Bern, the Swiss National Science Foundation, and the ESA PRODEX (PROgramme de Développement d'Expériences scientifiques) program. Work at Southwest Research Institute was supported by subcontract #1496541 from the Jet Propulsion Laboratory (JPL). Work at the Royal Belgian Institute for Space Aeronomy (BIRA-IASB) was supported by the Belgian Science Policy Office via PRODEX/ROSINA PRODEX Experiment Arrangement 90020. Work at the University of Michigan was funded by NASA under contract JPL-1266313.

References

- [1]** Balsiger, H., et al., 2007. ROSINA - Rosetta Orbiter Spectrometer for Ion and Neutral Analysis. *Space Science Reviews*, 128, 745-801.
- [2]** Hässig, M., et al., 2016. Isotopic composition of CO₂ in the coma of 67P/Churyumov-Gerasimenko measured with ROSINA/DFMS. *Astronomy & Astrophysics* (submitted).
- [3]** Glassmeier, K.-H., Boehnhardt, H., Koschny, D., Kührt, E., Richter, I., 2007. The Rosetta Mission: Flying Towards the Origin of the Solar System. *Space Science Reviews*, 128, 1-21.
- [4]** Lodders, K., 2003. Solar System Abundances and Condensation Temperatures of the Elements. *The Astrophysical Journal*, 591, 1220.
- [5]** McKeegan, K., Kallio, A., Heber, V., et al., 2011. The Oxygen Isotopic Composition of the Sun Inferred from Captured Solar Wind. *Science*, 332, 1528.
- [6]** Sakamoto, N., et al, 2007. Remnants of the Early Solar System Water Enriched in Heavy Oxygen Isotopes. *Science*, Vol 317, 13 July 2007.