

Diurnal gas cycle on 67P from ROSINA, rotation measurements, and cross-instrument identification of icy patches

Tobias Kramer (1,2), Matthias Lauter (1), Martin Rubin (3), and Kathrin Altwegg (3)

(1) Zuse Institute Berlin, Germany (kramer@zib.de, laeuter@zib.de), (2) Department of Physics, Harvard University, USA, (3) University of Bern, Physikalisches Institut, Switzerland

Abstract

We propose a consistent picture of the diurnal and seasonal gas emission [1] of 67P/C-G based on the

- in-situ gas density probed by COPS/DFMS, converted to surface emission rates [2, 3]
- the rotational period and axis orientation,
- the direct identification of icy patches on the nucleus.

Combining these three sources provides strong constraints on the surface activity of the nucleus and a cross-validation of multi-instrument data. Our model [2, 3] provides both, temporal and spatial resolution (110 m) across the entire mission.

References

- [1] Hansen, K.C., et al.: Evolution of water production of 67P/Churyumov-Gerasimenko: An empirical model and a multi-instrument study, MNRAS, 15, S491, 2016.
- [2] Kramer, T., Lauter, M., Rubin, M., and Altwegg, K.: Seasonal changes of the volatile density in the coma and on the surface of comet 67P/Churyumov-Gerasimenko, MNRAS, 469, S20–S28, 2017.
- [3] Lauter, M., Kramer, T., Rubin, M., and Altwegg, K.: Surface localization of gas sources on comet 67P/Churyumov-Gerasimenko based on DFMS/COPS data, arXiv:1804.06696, 2018.