

$^{16}\text{O}/^{18}\text{O}$ & $^{16}\text{O}/^{17}\text{O}$ in Water in the Coma of Comet 67P/Churyumov-Gerasimenko measured with the Rosetta/ROSINA Double Focusing Mass Spectrometer

I. Schroeder (1), K. Altwegg (1,2), H. Balsiger (1), J. J. Berthelier (3), J. De Keyser (4), B. Fiethe (5), S. A. Fuselier (6,7), S. Gasc (1), T. I. Gombosi (8), M. Rubin (1), T. Sémon (1), C.-Y. Tzou (1), S. F. Wampfler (2), P. Wurz (1,2)

- (1) Physikalisches Institut, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland
- (2) Center for Space and Habitability, University of Bern, Gesellschaftsstrasse 6, CH-3012 Bern, Switzerland
- (3) LATMOS, 4 Avenue de Neptune, F-94100 Saint-Maur, France
- (4) Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Ringlaan 3, B-1180, Brussels, Belgium
- (5) Institute of Computer and Network Engineering (IDA), TU Braunschweig, Hans-Sommer-Straße 66, D-38106 Braunschweig, Germany
- (6) Space Science Division, Southwest Research Institute, 6220 Culebra Road, San Antonio, TX 78228, USA
- (7) University of Texas at San Antonio, San Antonio, TX
- (8) Department of Atmospheric, Oceanic and Space Sciences, University of Michigan, 2455 Hayward, Ann Arbor, MI 48109, USA

Abstract

In an earlier study, the $^{16}\text{O} / ^{18}\text{O}$ ratio of H_2O in the coma of the comet 67P / Churyumov-Gerasimenko was found to be 445 ± 35 (Schroeder et al. 2018) [1]. The result was based upon in-situ measurements of $\text{H}_2^{16}\text{O} / \text{H}_2^{18}\text{O}$ and $^{16}\text{OH} / ^{18}\text{OH}$ performed with the ROSINA Double Focusing Mass Spectrometer (DFMS) on board the ESA spacecraft Rosetta and represented an 11% enrichment of ^{18}O as compared to the terrestrial value of 498.7 ± 0.1 measured by Baertschi (1976) [4]. This was consistent with leading self-shielding models, which had predicted primordial water in comets to be between 5 ~ 20 % more enriched in ^{18}O than terrestrial water [6]. The $^{16}\text{O} / ^{17}\text{O}$ ratio, however, could not be easily determined at the time due to the low signal from the H_2^{17}O peak and its overlap with the much larger HDO peak.

In contrast, CO_2 in the coma of 67P has a $^{16}\text{O} / ^{18}\text{O}$ ratio of 494 ± 8 , as previously determined by Hässig et al. (2016) [2] using the same instrument, which is consistent within 1σ uncertainty with the terrestrial value. The solar wind measured by McKeegan et al. (2011) [5], on the other hand, exhibited ^{18}O -depletion with a $^{16}\text{O} / ^{18}\text{O}$ ratio of 530 ± 2 .

In this follow-up study, we separate the H_2^{17}O peak from that of HDO by manually fitting them with equal-width Gaussians and thus obtain an estimate for the $^{16}\text{O} / ^{17}\text{O}$ ratio of H_2O in 67P's coma from several DFMS mass spectra that were acquired during 67P's outbound equinox in March 2016 and which were chosen for their strong signal.

Acknowledgements

ROSINA would not have produced such outstanding results without the work of the many engineers, technicians and scientists involved in the mission, in the Rosetta spacecraft and in the ROSINA instrument team over the last 20 years, whose contributions are gratefully acknowledged. Rosetta is a European Space Agency (ESA) mission with contributions from its member states and NASA. We acknowledge herewith the work of the whole ESA Rosetta team.

Work at University of Bern was funded by the State of Bern, the Swiss National Science Foundation, and the ESA PRODEX (PROgramme de Développement d'Expériences scientifiques) program. Work at Southwest Research Institute was supported by subcontract #1496541 from the Jet Propulsion Laboratory (JPL). Work at the Royal Belgian Institute for Space Aeronomy (BIRA-IASB) was supported by the Belgian Science Policy Office via PRODEX/ROSINA PRODEX Experiment Arrangement 90020. Work at the University of Michigan was funded by NASA under contract JPL-1266313.

References

- [1] Schroeder I. et al., 2018, The $^{16}\text{O} / ^{18}\text{O}$ Ratio in Water in the Coma of Comet 67P / Churyumov-Gerasimenko measured with the Rosetta / ROSINA Double Focusing Mass Spectrometer. A&A Rosetta Special Issue.
- [2] Hässig, M., et al., 2016. Isotopic composition of CO₂ in the coma of 67P / Churyumov-Gerasimenko measured with ROSINA / DFMS. Astronomy & Astrophysics (submitted).
- [3] Glassmeier, K.-H., Boehnhardt, H., Koschny, D., Kührt, E., Richter, I., 2007. The Rosetta Mission: Flying Towards the Origin of the Solar System. Space Science Reviews, 128, 1-21.
- [4] Baertschi P., 1976, Absolute ^{18}O content of standard mean ocean water. Earth and Planetary Science Letters, Volume 31, Issue 3, 341-344.
- [5] McKeegan, K., Kallio, A., Heber, V., et al., 2011. The Oxygen Isotopic Composition of the Sun Inferred from Captured Solar Wind. Science, 332, 1528.
- [6] Sakamoto, N., et al, 2007. Remnants of the Early Solar System Water Enriched in Heavy Oxygen Isotopes. Science, Vol 317, 13 July 2007.