

## Detection of organic material on the nucleus of comet 67P: insight into their origin from IR spectral features.

A. Raponi<sup>1</sup>, M. Ciarniello<sup>1</sup>, G. Filacchione<sup>1</sup>, F. Capaccioni<sup>1</sup>, M.C. De Sanctis<sup>1</sup>, L. V. Moroz<sup>3,4</sup>, V. Vinogradoff<sup>1</sup>, F. Tosi<sup>1</sup>, G. Arnold<sup>3</sup>, E. Quirico<sup>2</sup>, V. Mennella<sup>7</sup>, P. Beck<sup>2</sup>, C. Pilorget<sup>5</sup>, A. Pommerol<sup>6</sup>, S. Schröder<sup>3</sup>, D. Kappel<sup>3</sup>, I. Istiqomah<sup>2</sup>, B. Rousseau<sup>2</sup>, O. Poch<sup>2</sup>

<sup>1</sup>IAPS-INAF, Rome, Italy (andrea.raponi@iaps.inaf.it); <sup>2</sup>Institut de Planétologie et d'Astrophysique de Grenoble, Grenoble, France; <sup>3</sup>German Aerospace Center (DLR), Berlin, Germany; <sup>4</sup>Institute of Earth and Environmental Science, Univ. of Potsdam, Germany; <sup>5</sup>IAS Institut d'Astrophysique Spatiale, Orsay Cedex, France; <sup>6</sup>University of Bern, Bern, Switzerland; <sup>7</sup>OACN-INAF, Naples, Italy

We revised the in-flight calibration of the VIRTIS-M-IR instrument onboard Rosetta spacecraft with the aim to improve: a) the detection of low-contrast spectral features and b) the radiometric accuracy. Furthermore, we have modeled the thermal emission to remove the nucleus contribution at wavelengths in excess of 3.0  $\mu\text{m}$  and we have also modeled and removed, by means of the Hapke [1] model, the contribution of water ice absorption to isolate the organic features within the spectral region 2.9 – 3.5  $\mu\text{m}$ . We identified small features centered at: 3.28  $\mu\text{m}$ , which are consistent with aromatic C-H stretching; 3.38, 3.42, 3.48  $\mu\text{m}$  which can be attributed to aliphatic ( $\text{CH}_3$  asymmetric,  $\text{CH}_2$  asymmetric, and  $\text{CH}_3$  symmetric stretching, respectively) [2]; other small features are present at 2.85, 3.0, 3.1  $\mu\text{m}$ . They can be attributed to C-H overtones, OH-stretches, and/or N-H stretching, according to spectral comparison with analog materials. Finally, the organic features on the spectrum of the comet present similarities with observations of interstellar diffuse material [3], and protoplanetary nebulae [4]. Their significance in the framework of the comet origin and evolution will be discussed.

**References.** [1] Hapke B., 2012, Theory of Reflectance and Emittance Spectroscopy, 2nd edn. Cambridge Univ. Press, Cambridge; [2] L. V. Moroz et al. Icarus, 134, pp. 253; [3] Dartois E. et al., 2004, A&A, 423, 549; [4] Sandford S. A. et al., 2013, ApJS, 205.

**Acknowledgements.** We thank the following institutions and agencies for support of this work: Italian Space Agency (ASI, Italy) Centre National d'Etudes Spatiales (CNES, France), DLR (Germany). This work takes advantage of the collaboration of the ISSI international team “Comet 67P/Churyumov-Gerasimenko Surface Composition as a Playground for Radiative Transfer Modeling and Laboratory Measurements”, number 397.